

25 PCT 1994 14.07.00

LITERATUR

5254/09/00



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b>  <b>C12P 21/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/07287</b>  <b>(43) International Publication Date:</b> 15 April 1993 (15.04.93)
<b>(21) International Application Number:</b> PCT/US92/08518 <b>(22) International Filing Date:</b> 7 October 1992 (07.10.92)  <b>(30) Priority data:</b> 775,136 11 October 1991 (11.10.91) US  <b>(71) Applicant:</b> PROMEGA CORPORATION [US/US]; 2800 Woods Hollow Road, Madison, WI 53711 (US).  <b>(72) Inventors:</b> THOMPSON, David, V. ; 611 West Dean Avenue, Monona, WI 53716 (US). VAN OOSBREE, Thomas, R. ; 2123 Commonwealth, Madison, WI 53705 (US).  <b>(74) Agent:</b> GEMIGNANI, Joseph, A.; Michael, Best & Friedrich, 100 East Wisconsin Avenue, Milwaukee, WI 53202-4108 (US).		<b>(81) Designated States:</b> AU, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE).  <b>Published</b> <i>With international search report</i> <i>With amended claims and statement</i>
<b>(54) Title:</b> COUPLED TRANSCRIPTION AND TRANSLATION IN EUKARYOTIC CELL-FREE EXTRACT  <b>(57) Abstract</b>  A method for coupling transcription and translation from DNA using a solution comprising a eukaryotic cell-free extract comprising adding a sufficient amount of a magnesium compound to the extract to raise the magnesium concentration to a level where RNA is transcribed from DNA and RNA translates into protein.		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SK	Slovak Republic
CI	Côte d'Ivoire	LK	Sri Lanka	SN	Senegal
CM	Cameroon	LU	Luxembourg	SU	Soviet Union
CS	Czechoslovakia	MC	Monaco	TD	Chad
CZ	Czech Republic	MG	Madagascar	TG	Togo
DE	Germany	ML	Mali	UA	Ukraine
DK	Denmark	MN	Mongolia	US	United States of America
ES	Spain			VN	Viet Nam
FI	Finland				

COUPLED TRANSCRIPTION AND TRANSLATION  
IN EUKARYOTIC CELL- FREE EXTRACT

BACKGROUND OF THE INVENTION

This invention relates generally to molecular  
5 biology, and more particularly, to a new method allowing  
coupling of the transcription of RNA from a template DNA  
and the translation of the RNA in eukaryotic cellular  
lysates or other extracts.

The steps involved in the transcription and  
10 translation (expression) of genes in cells are very  
complex and are not yet completely understood. There is  
a basic pattern that must be followed, however, for  
protein to be produced from DNA. The DNA is first  
transcribed into RNA, and then the RNA is translated by  
15 the interaction of various cellular components into  
protein. In prokaryotic cells (bacteria) transcription  
and translation are "coupled", meaning that RNA is  
translated into protein during the time that it is being  
transcribed from the DNA. In eukaryotic cells (animals,  
20 plants) the two activities are separate, making the  
overall process much more complicated. DNA is  
transcribed into RNA inside the nucleus of the cell, but  
the RNA is further processed into mRNA and then  
transported outside the nucleus to the cytoplasm where  
25 it is translated into protein.

The ability of molecular biologists to isolate and  
clone genes, as well as their ability to isolate  
particular mRNAs or "messages" from cells, has brought  
about the need for systems which can be used to express  
30 these genes or messages. The expression of a gene is

important in the overall understanding of its function and regulation. Methods are now available for rapid expression of proteins, making it possible to manipulate genes and then study the effect of the manipulations on their function. The amount of protein to be produced, whether the gene is prokaryotic or eukaryotic and the relative merits of an in vitro or an in vivo system, are some of the factors considered by researchers when selecting an expression system. The choice of a system is influenced by the gene being studied. For the most part, a prokaryotic gene is expressed best in a prokaryotic system, and a eukaryotic gene is more efficiently and accurately expressed in a eukaryotic system. This is because of the many regulatory sequences and promoters that are recognized more efficiently in a like system. The expression of genes can be achieved in both in vivo and in vitro systems.

In vivo transcription systems using prokaryotic or eukaryotic cells are available, however, these systems are difficult to work with since intact cells are used. In vitro systems, on the other hand, are made from cell-free extracts produced from prokaryotic or eukaryotic cells that contain all the necessary components to translate DNA or RNA into protein. Cell-free extracts can be prepared from prokaryotic cells such as E. coli and from eukaryotic cells such as rabbit reticulocytes and wheat germ. Cell-free systems are very popular because there are standard protocols available for their

preparation and because they are commercially available from a number of sources.

E. coli S30 cell-free extracts were first described by Zubay, G. (1973, Ann. Rev. Genet. Vol 7, p. 267).

5 These can be used when the gene to be expressed has been cloned into a vector containing the appropriate prokaryotic regulatory sequences, such as a promoter and ribosome binding site. Prokaryotic E. coli cell-free systems are considered "coupled" because transcription  
10 and translation occur simultaneously after the addition of DNA to the extract. The use of RNA as a template in E. coli extracts results in protein production but such a reaction is not coupled. Rabbit reticulocyte lysates and wheat germ extracts are used preferably for the  
15 expression of eukaryotic genes or mRNA. Both systems require the use of RNA as the template for protein translation because, as previously mentioned, eukaryotic systems are not coupled.

Rabbit reticulocyte lysate was described by Pelham, H.R.B. and Jackson, R.J. (1976, Eur. J. Biochem. Vol. 131, p. 289). This expression system is probably the most widely used cell-free system for in vitro translation, and is used in the identification of mRNA species, the characterization of their products, the  
25 investigation of transcriptional and translational control. Processing events, such as signal peptide cleavage and core glycosylation, are examined by adding canine microsomal membranes to a standard translation reaction (Walter, P. and Biobel, G. (1983) Meth.

Enzymol. 96, 50). Rabbit reticulocyte lysate also contains a variety of post-translational processing activities, including acetylation, isoprenylation, proteolysis and some phosphorylation activity (Glass, 5 C.A. and Pollard, K.M. (1990) Promega Notes 26).

Wheat germ extract was described by Roberts, B.C. and Paterson, B.M. (1973, Proc. Natl. Acad. Sci. USA, Vol. 70, P. 2330). Cell-free extracts of wheat germ support the translation in vitro of a wide variety of 10 viral and other prokaryotic RNAs, as well as eukaryotic mRNAs. (Anderson, C., et al. (1983) Meth. Enzymol. 101, 635). Generally, it is found necessary to include a ribonuclease inhibitor in the reaction mix of a wheat germ translation system, as ribonuclease activities in 15 wheat germ extract are present.

RNA for translational studies is obtained by either isolating mRNA or by making in vitro RNA transcripts from DNA that has been cloned into a vector containing an RNA polymerase promoter. The first method isolates 20 mRNA or "message" directly from cells.

The second obtains RNA for in vitro translation by in vitro transcription. In vitro transcription of cloned DNA behind phage polymerase promoters was described by Krieg, P. and Melton, D (1984, Nucl. Acids 25 Res., Vol. 12, p. 7057). This has become a standard method for obtaining RNA from cloned genes for use in in vitro translation reactions. This method requires that the DNA or gene of interest be cloned into a vector containing a promoter for one of the following RNA

- 5 -

polymerases, SP6, T7 or T3. The vector is then linearized at the 3' end of the cloned gene using a restriction enzyme, followed by an in vitro transcription reaction to make RNA transcripts. A number of vectors containing the SP6, T7 and T3 RNA polymerase promoters are commercially available and are widely used for cloning DNA.

In any case, the process of obtaining RNA transcripts for use in rabbit reticulocyte lysate or wheat germ systems introduces a variable which can affect the efficiency of the translation reaction. Extra care must always be taken when working with RNA as it is easily degraded by ribonucleases. DNA templates are much more stable.

After rabbit reticulocyte lysate and wheat germ extract were developed as cell-free translation systems, attempts were made to couple transcription and translation. One system that was developed was a "linked" transcription and translation system (Roberts, B.E., et al. (1975), Proc. Natl. Acad. Sci. USA, Vol 72, 1922-1926). This system involved the use of wheat germ extract and looked at transcription and translation of SV40 viral DNA using E. coli RNA polymerase. In this system transcription occurs in 15 minute incubation step just prior to the addition of the wheat germ extract. The steps are separated because of incompatibility between the buffer conditions necessary for transcription and those necessary for translation, and also because of the different temperature requirements

for both processes. This system has a number of drawbacks. One is the lack of control over which protein product is produced, as a number of different proteins are synthesized simultaneously from the same SV40 DNA template. Although the authors of that study indicated that a coupled system had been developed no data for a coupled system was shown.

Another system was developed by Pelham, H.R.B, et al. (1978), Eur. J. Biochem., Vol. 82, 199-209, where coupled transcription and translation occurred after the introduction of vaccinia viral core particles into rabbit reticulocyte lysate. The production of vaccinia proteins from the viral DNA was presumably due to transcription by the endogenous vaccinia RNA polymerase and subsequent translation by the lysate. This system was limited by the fact that only vaccinia proteins would be produced while exogenous DNA from sources other than vaccinia would not be recognized by the RNA polymerase and therefore no transcription or translation could occur. Viral core particles had to be isolated, and the authors were unable to exclusively produce a single protein.

Work has also been described using "continuous" cell-free in vitro translation systems with the emphasis on large scale production of protein. Continuous systems are very different than the more common batch type, or static, in vitro translation reactions which occur in a contained reaction volume. Continuous translation involves a bioreactor (such as an Amicon 8MC



ultrafiltration unit) in which large scale reactions are set up and protein is "continually" translated over extended periods of time. The reaction requires that a buffer be fed into the reaction as it progresses, and also requires that the products of translation be removed from the reaction filter unit. This type of system works well with E. coli S30 extract and wheat germ extract when RNA template is introduced. See Spirin, et al. (1988) Science, Vol 242, 1162-1164. The system also works using RNA templates in rabbit reticulocyte lysate. See Ryabova, et al. (1989) Nucl. Acid Res., Vol. 17, No. 11, 4412. The system is also known to work well with DNA templates in E. coli S30 extracts. See Baranov, et al. (1989) Gene, Vol 84, 463-466. PCT publication WO9102076 discloses continuous cell-free translation from DNA templates using eukaryotic lysates.

Continuous reactions are performed variously from tens of hours up to around one hundred hours, and require a substantial investment of time and resources to set up and run. Translation in a "continuous" system is also directed towards producing large amounts of protein, and differs substantially from standard (static) in vitro translation reactions. Static reactions can be run in a small reaction volume, typically measured in microliters, and are often completed in one to two hours. A static translation reaction is not directed towards producing preparative amounts (milligrams) of proteins. A static reaction is

generally used to produce protein for investigative applications, such as the identification and characterization of mRNA species, or studies of transcriptional or translation control. None of the  
5 rabbit reticulocyte or wheat germ systems currently known for in vitro translation provide for coupled transcription and translation in a static reaction mixture.

#### SUMMARY OF THE INVENTION

10 It is an object of the invention, therefore, to provide a simple method for producing protein from a template DNA in a standard in vitro translation reaction utilizing a eukaryotic cellular lysate.

A more specific object of the invention is to  
15 provide such a method which can be used to couple transcription and translation of a single protein coded by the DNA template.

Another object is to provide a process for enhanced  
20 in vitro protein production from DNA templates through coupled transcription and translation reactions using eukaryotic cell-free lysates.

A further object is to provide a method for manufacturing eukaryotic cellular lysates for use in coupled transcription and translation experiments.

25 A still further object is to provide a method for producing such a lysate preparation which has one or more of the components or reagents necessary for coupled transcription and translation.

It is another object to provide such a preparation for use in coupled transcription and translation experiments.

A further object of the invention is to provide a  
5 kit having a prepared rabbit reticulocyte lysate or wheat germ extract and including other components or reagents necessary for producing a reaction mixture for coupling transcription and translation from a DNA template.

10 A more specific object is to provide a kit having as at least one component a prepared rabbit reticulocyte lysate or wheat germ extract preparation which has one or more of the reagents necessary for coupling transcription and translation.

15 For the achievement of these and other objects, this invention provides a method for coupling transcription and translation in eukaryotic cellular lysate. The method comprises adding a sufficient amount of a magnesium compound to the lysate to raise the final  
20 magnesium concentration thereof to a level where RNA is transcribed from DNA and RNA translates into protein.

The invention also provides a process for producing protein from a DNA template having a specific polymerase promoter sequence through coupled transcription and  
25 translation in a solution. This is accomplished by preparing a standard preparation of a eukaryotic cellular lysate, modifying the lysate with sufficient concentrations of ribonucleotide triphosphates, amino acids and the polymerase corresponding to the promoter

sequence of the template DNA to support transcription and translation, and adding a sufficient amount of a magnesium compound to the solution to raise the final magnesium concentration to a level where RNA is transcribed from DNA and RNA translates into protein. Only that protein corresponding to the gene adjacent the promoter sequence in the DNA will be produced.

The invention additionally provides a modified eukaryotic cellular lysate prepared in accordance with the method whereby the final magnesium concentration is raised to a level where RNA is transcribed from DNA and RNA translates into protein. The lysate can have its magnesium concentration so adjusted during manufacture. The invention also provides a kit that comprises a container containing prepared eukaryotic cellular lysate for coupled transcription and translation. In addition to having an adjusted magnesium concentration, this lysate can contain a variety of additional components required for coupled transcription and translation. One of the additional components that can be included is an RNA polymerase. Others are various buffers or salts or other components or reagents for optimizing coupled transcription and translation reaction conditions, such as spermidine, which is known to stimulate the efficiency of chain elongation.

The coupling of transcription and translation in rabbit reticulocyte or wheat germ systems significantly increases protein production when compared to standard in vitro translation with RNA templates, and the coupled

transcription and translation has been shown to work for a variety of different proteins over a wide range of molecular weight sizes.

The gene or DNA to be translated is preferably  
5 cloned behind an RNA polymerase promoter such as the SP6, T7, or T3 promoters, or any RNA polymerase promoter for which a polymerase is available. The DNA template can be introduced into the coupled transcription and translation reaction in the form of a closed circular  
10 plasmid DNA or as a linearized plasmid DNA. RNA polymerase promoter sequences can also be attached to specific DNA fragments by the thermocycling amplification process, and these linear DNA fragments can be used in coupled transcription and translation  
15 reactions.

Standard in vitro translation reactions require very little investment of time and resources and are a well established method for expressing protein qualitatively. By coupling transcription and  
20 translation in standard in vitro translation reactions using rabbit reticulocyte lysate or wheat germ extract the need for a separate in vitro transcription reaction is eliminated and protein production is significantly enhanced. Thus, coupled transcription and translation  
25 using rabbit reticulocyte lysate or wheat germ extract provides a significant improvement over current standard in vitro translation reactions. An unexpected benefit is that the production of the desired protein product is increased dramatically (several fold) over that which is

-12-

seen by the addition of RNA to standard in vitro translation reactions. In addition, there is no need for a separate capping reaction with coupled transcription and translation.

5 Another advantage is that protein production can be achieved from a cloned or amplified DNA template containing a specific RNA polymerase promoter. In this manner RNA can be directed to be synthesized from a single gene downstream of the promoter, and only the  
10 proteins translated from that RNA are produced. If that RNA directs only one protein, then only one protein is produced through the reaction. Plasmid vectors for such use are well known in the art and are available from several sources, allowing the researcher to produce  
15 protein from any cloned gene at will.

Other features and advantages of the invention will become apparent to those of ordinary skill in the art upon review of the following detailed description and claims.

20 DESCRIPTION OF THE PREFERRED EMBODIMENT

For purposes of the invention, any eukaryotic cellular lysate can be used, and a number of conventional techniques exist for their preparation. Eukaryotic cell-free lysates are preferred expression  
25 systems for many reasons, at least partially because they retain a variety of post-translational processing activities. With the addition of canine microsomal membranes processing events, such as signal peptide

- 13 -

cleavage and core glycosylation, can be examined.

Eukaryotic cellular lysates also support the translation in vitro of a wide variety of viral and other prokaryotic RNAs, as well as eukaryotic mRNAs.

5           While other eukaryotic systems are suitable, rabbit reticulocyte lysate and wheat germ extract are preferred. These eukaryotic lysates are popular with researchers, and are widely available. In a preferred embodiment, rabbit reticulocyte lysate is prepared by a  
10           method described by Pelham, H. and Jackson, R.J. (1976, Eur. J. Biochem. 67, 247-256) and modified according to the manufacturing protocol L415/L416, Promega Corp., Madison, WI. Reticulocyte lysate is prepared from New Zealand White rabbits injected with phenylhydrazine,  
15           which ensures reliable and consistent reticulocyte production in each lot. The reticulocytes are purified to remove contaminating cells which could otherwise alter the translational properties of the final extract. After the reticulocytes are lysed, the extract is  
20           treated with micrococcal nuclease and  $\text{CaCl}_2$ , to destroy endogenous mRNA and thus reduce background translation to a minimum. EGTA is then added to chelate the  $\text{CaCl}_2$  and thereby inactivate the nuclease.

          The lysate contains cellular components necessary  
25           for protein synthesis. These include tRNAs, rRNAs, amino acids and initiation, elongation, and termination factors. The lysate is further optimized for mRNA translation by adding an energy generating system, consisting of pre-tested phosphocreatine kinase and

-14-

phosphocreatine. Also added are a mixture of tRNAs to expand the range of mRNAs which can be translated, and hemin to prevent inhibition of initiation. Hemin is included in the reticulocyte lysate because it is a suppressor of an inhibitor of the initiation factor eIF2 $\alpha$ . In the absence of hemin, protein synthesis in reticulocyte lysates ceases after a short period of incubation (Jackson, R. and Hunt, T. 1983 Meth. In Enzymol. 96, 50). Potassium acetate and magnesium acetate are added at a level recommended for the translation of most mRNA species. This is the standard rabbit reticulocyte lysate used for in vitro translations. The final magnesium concentrations for standard rabbit reticulocyte lysate are typically in the range of about 4.2 to 5.0 mM, which is used at a 50% ratio in coupled transcription and translation reactions.

---

Final Concentration Contributions of Added Components  
in a Rabbit Reticulocyte Lysate Translation Reaction  
Which is 50% Lysate

---

25	Creatine phosphate	7 mM
	Creatine phosphokinase	35 $\mu$ g/ml
	DTT	1.4 mM
	Calf liver tRNA	35 $\mu$ g/ml
	Potassium acetate	56 mM
30	Magnesium acetate	350 $\mu$ M
	Hemin	14.3 $\mu$ M

---

Another preferred embodiment utilizes wheat germ extract. This can be prepared by a method described by Roberts, B.E. and Paterson, B.M. (1973), Proc. Natl. Acad. Sci. USA Vol. 70, No. 8, pp. 2330-2334), following



the modifications described by Anderson, C.W., et al. (1983, Meth. Enzymol. Vol.101, p. 635) and modified as in the manufacturing protocol L418, Promega Corp. Madison, WI. Generally, wheat germ extract is prepared

5 by grinding wheat germ in an extraction buffer, followed by centrifugation to remove cell debris. The supernatant is then separated by chromatography from endogenous amino acids and plant pigments that are inhibitory to translation. The extract is also treated

10 with micrococcal nuclease to destroy endogenous mRNA, to reduce background translation to a minimum. The extract contains the cellular components necessary for protein synthesis, such as tRNA, rRNA and initiation, elongation, and termination factors. The extract is

15 further optimized by the addition of an energy generating system consisting of phosphocreatine kinase and phosphocreatine, and magnesium acetate is added at a level recommended for the translation of most mRNA species. The final magnesium concentration for standard

20 wheat germ extract is typically in the range of about 6.0 mM to 7.5 mM.

---

Final Concentration Contributions of Added  
Components in a Wheat Germ Extract Translation  
Reaction Which is 50% Extract

---

5.		
	Creatine phosphate	10 mM
	Creatine phosphokinase	50 µg/ml
	DTT	5 mM
	Calf liver tRNA	50 µg/ml
10	Magnesium acetate	3.0 - 3.75 mM
	Potassium acetate	50 mM
	Spermidine	0.5 mM
	ATP	1.2 mM
	GTP	0.1 mM
15	HEPES (pH 7.6)	12 mM

---

For coupled transcription and translation the magnesium concentration of the eukaryotic cellular lysate must be adjusted by an additional magnesium compound, preferably a salt. Preferred salts include magnesium chloride and magnesium acetate. The addition of a buffering agent can be used in the solution to stabilize the pH, although this isn't necessary. For coupling transcription and translation, a sufficient amount of magnesium chloride or acetate is added to the lysate to raise the final magnesium concentration to a level where RNA is transcribed from DNA and RNA translates into protein. This level will vary depending upon the lysate used.

The simple addition of a prokaryotic RNA polymerase and ribonucleotide triphosphates to a standard rabbit reticulocyte lysate or wheat germ extract does not allow in vitro protein production from DNA templates. The particular adjustments of the salt concentrations in the system that will be described, however, allow protein production by creating conditions within the lysate

which permit both transcription of DNA into RNA and translation of the RNA into protein.

Magnesium is known to be important for optimizing translation, as it enhances the stability of assembled  
5 ribosomes and functions in their binding together during translation. Magnesium also appears to play a role in facilitating polymerase binding. Potassium is important as well for optimizing translation, but unlike the case for magnesium, for coupled transcription and translation  
10 the concentration of potassium ions does not need to be altered beyond standard translation preparation levels.

Potassium and magnesium are in the standard rabbit lysate. The levels are partially from the endogenous  
lysate levels, and partially from the additions made in  
15 the preparation of the lysate, as are done for translation lysates. Lysate is diluted somewhat in manufacture, and the prepared lysate is then only used at 50% for coupled transcription and translation reactions.

20 As the magnesium concentration should be adjusted to within a rather narrow optimal range, it is preferred that the lysate magnesium levels be measured directly through the use of a magnesium assay, prior to the addition of extra magnesium, so that the amount of  
25 magnesium in a reaction can be standardized from one batch of lysate to the next. The Lancer "Magnesium Rapid Stat Diagnostic Kit" (Oxford Lab Ware Division, Sherwood Medical Co., St. Louis, MO), is one such assay which can accurately measure the magnesium levels in

-18-

biological fluid. Once the magnesium ion concentration for a given batch of lysate is known then additional magnesium, for instance in the form of a concentrated magnesium salt solution, can be added in a known manner to bring the magnesium concentration of the lysate to within the optimal range, or, in the case of a modified lysate preparation to be used as one-half of a reaction mixture, to within twice the optimal range.

Thus, it has been found that when the final magnesium concentration of rabbit reticulocyte lysate is adjusted, such as by adding a concentrated solution of magnesium chloride or acetate, to a concentration greater than 2.5 mM but less than 3.5 mM, preferably between 2.6 mM and 3.0 mM, coupled transcription and translation occurs. For coupling transcription and translation using wheat germ extract, a final concentration of magnesium chloride or acetate greater than about 3.0 mM but less than about 5.25 mM produces protein, preferably adjusted to approximately 4.0 mM to 4.75 mM.

Reaction conditions for coupled transcription and translation must include the addition of ribonucleotide triphosphates (ATP, GTP, CTP, UTP) and amino acids, for rabbit reticulocyte lysate to final concentrations of 0.4 mM each, and 20  $\mu$ M each respectively. Reaction conditions for wheat germ extract are modified by the addition of ribonucleotide triphosphates to final concentrations of 0.4 mM for CTP and UTP, 0.5 mM for GTP and 1.6 mM for ATP, while amino acids are added to a

final concentration of 20-80  $\mu$ M. If a radiolabeled amino acid is used in the coupled reaction, such as  $^{35}$ S methionine or  $^3$ H leucine, then the corresponding amino acid is left out of the amino acid mix. An RNA  
5 polymerase, either SP6, T7, or T3, is then added, preferably to a final concentration of about 80-160 units per 50  $\mu$ l reaction. The DNA template with the gene to be translated is added at a concentration of 1  $\mu$ g, and the reaction volume is adjusted to 50  $\mu$ l with  
10 the addition of water. The reaction is then incubated at 30°C for 1-2 hours.

Although potassium is added to the reaction mixture in the preferred embodiment, in contrast to magnesium additional potassium does not greatly increase protein  
15 production, but only offers a slight improvement when proper magnesium levels are already present. Potassium acetate is added to an optimal final concentration of about 59 mM. Although potassium chloride or acetate can be added, because of the greater amounts added than is  
20 the case for magnesium, potassium acetate is preferred. The standard translation lysate level can be used, a concentration of approximately 56 mM, while spermidine is added to give a final concentration of about 0.2 mM. The final concentration of potassium chloride or acetate  
25 is also an estimation based on the amount of this component in standard lysate, but it must be recognized that this concentration, as well as the magnesium concentration, can vary slightly due to endogenous components.

Additional components can be added to the lysate as desired for improving the efficiency or stability of the coupled transcription and translation reaction. One common addition to coupled transcription and translation reactions is an amount of a polyamine sufficient to stimulate the efficiency of chain elongation. Although not absolutely necessary, for coupled transcription and translation a final concentration of spermidine in the mixture of about 0.2 mM is optimal, in that an increase of protein production is observed with this concentration. Polyamines affect optimal magnesium levels as well, and are known to lower the effective magnesium concentration for translation reactions somewhat. It appears that the polyamines may substitute for magnesium at some level, and thus would play a role in the optimization of magnesium requirements, possibly even permitting some lowering of optimal magnesium levels for coupled transcription and translation.

Optimal magnesium concentrations in the in vitro environment are affected by other conditions and considerations, too. As the ribonucleotide triphosphate concentration goes up, for instance, there is a concomitant increase in the optimal magnesium concentration, as the ribonucleotide triphosphates tend to associate, or chelate, with magnesium in solution. Thus, when the ribonucleotide triphosphate concentrations cited for the above reactions are increased to 0.6 mM, the production of protein from coupled transcription and translation reactions is

greatly reduced. The optimal concentration of magnesium also varies with the type of cellular lysate, whether using wheat germ extract or rabbit reticulocyte lysate. The amount of magnesium required to be added to achieve  
5 optimal levels will vary with the concentration of the lysate used in the reaction mixture, as increasing the concentration of the lysate will increase the contribution of magnesium from the lysate itself.

Due to the large number of components in a lysate  
10 mixture, it can not be said with certainty whether it is the protein translation from RNA, the transcription of RNA from DNA or both that is adversely affected at other than optimal salt conditions. The observation is the same in any case, that detectible levels of protein are  
15 not produced in the reaction. With a small adjustment to the magnesium concentration, possibly adjusted by the polyamine concentration and watching that ribonucleotide triphosphate concentration levels do not become a problem, coupled transcription and translation is  
20 observed, but only through a relatively small range.

Dithiothreitol (DTT) is preferably added to the translation mixture. When included in coupled transcription and translation reactions, DTT is preferably added to a final concentration of about 1.45  
25 mM. Optimal DTT is about 5.1 mM for wheat germ. Also, a ribonuclease inhibitor, such as RNasin, can be added to the lysate, to effectively inactivate endogenous ribonucleases. Concentrations of 40 units per 50  $\mu$ l reaction have been shown to help prolong the reaction.

it is not an absolute requirement for coupled transcription and translation in rabbit reticulocyte lysate, but RNasin is required for coupled transcription and translation using wheat germ extract. The omission  
5 of RNasin from the latter results in no protein translation, as there are active ribonucleases present in the lysate.

The preferred coupled transcription and translation concentrations for rabbit reticulocyte lysate, of  
10 potassium chloride or acetate, magnesium chloride or acetate and spermidine, can be achieved by the addition of 2.5  $\mu$ l of an optimized tris/acetic acid buffer (1xTA buffer = 33 mM Tris/acetic acid pH 7.8, 65 mM potassium acetate, 10 mM magnesium acetate, 4 mM spermidine, 1 mM  
15 DTT). When added to a standard 50  $\mu$ l in vitro translation reaction, this buffer raises the magnesium level in the lysate by 0.5 mM overall.

Rabbit reticulocyte lysate can be modified during manufacture. The lysate is used at a 50% concentration  
20 (25  $\mu$ l lysate per 50  $\mu$ l reaction, typically), and so a preferred modification involves adjusting the potassium acetate concentration to 118 mM, the magnesium acetate to about 5.2 mM to 6.0 mM, and spermidine to 0.4 mM. This gives optimal final concentrations of 59 mM  
25 potassium acetate, 2.6 mM to 3.0 mM magnesium acetate and 0.2 mM spermidine when 50% lysate is used in a coupled transcription and translation reaction. The lysate can be further modified by the addition of one of the RNA polymerases (SP6, T7, or T3) to the lysate,



-23-

preferably at a concentration of 80-160 units per 50  $\mu$ l reaction. Such a modified lysate can be stored frozen until needed.

Spermidine is also preferably added to wheat germ  
5 extract, optimally to a final concentration of about 0.9 mM. Potassium acetate is preferably added to a final concentration of approximately 56.5 mM. These concentrations can be achieved by the addition of 5.0  $\mu$ l of 1xTA buffer to a 50  $\mu$ l in vitro translation using  
10 standard wheat germ extract, although these concentrations are based on estimations of the amounts of these components in standard wheat germ extract and may vary slightly due to endogenous components in the lysate itself.

15 Wheat germ extract can also be modified during the manufacturing process so that the final concentrations of potassium acetate, magnesium acetate and spermidine will equal those described in the reaction conditions above for wheat germ when the lysate is used at a 50%  
20 concentration. Wheat germ extract can be further modified during manufacture by the addition of one of the RNA polymerases to a final concentration of 80-160 units per reaction. Modified extract is stored frozen until it is used.

25 Leaving magnesium concentrations at levels present in the standard lysate, production of protein does not occur. The addition of all of the other components of the 1xTA buffer to the reaction mixture, only without the magnesium, results in a reaction that produces no

-24-

protein. On the other hand, addition of excess magnesium will cease translation altogether. For instance, in a similar reaction mixture utilizing rabbit reticulocyte lysate, but having final magnesium concentrations of about 3.5 mM, translation of protein again ceases. This upper limit will presumably vary slightly depending upon other parameters such as potassium and spermidine concentrations, as well as ribonucleotide triphosphate concentrations both of which are known to vary the optimal magnesium concentration for translation of RNA into proteins.

Both modified wheat germ extract and modified rabbit reticulocyte lysate can be included as part of a kit for facilitating the set up of coupled transcription and translation reactions. Such a kit improves the convenience to the researcher, as the eukaryotic cellular lysate comes prepared and ready for use. In addition to the rabbit reticulocyte lysate or wheat germ extract, such a kit can include the components, reagents, including enzymes, and buffers necessary to perform coupled transcription and translation upon the introduction of a DNA template. The lysate can be standard, or can be of the type where the adjustments to its salt concentrations have already been made during manufacture, or additionally where one or more of the components, reagents or buffers necessary for coupled transcription and translation have been included.

The amount of protein produced in a coupled in vitro transcription and translation can be measured in

various fashions. One method relies on the availability of an assay which measures the activity of the particular protein being translated. An example of an assay for measuring protein activity is the luciferase assay system described in Technical Bulletin 097, Promega Corp., Madison, WI. These assays measure the amount of functionally active protein produced from the coupled in vitro transcription and translation reaction. Activity assays will not measure full length protein that is inactive due to improper protein folding or lack of other post translational modifications necessary for protein activity.

Another method of measuring the amount of protein produced in coupled in vitro transcription and translation reactions is to perform the reactions using a known quantity of radiolabeled amino acid such as <sup>35</sup>S methionine or <sup>3</sup>H leucine and subsequently measuring the amount of radiolabeled amino acid incorporated into the newly translated protein. For a description of this method see the in vitro Translation Technical Manual, Promega Corp., Madison, WI. Incorporation assays will measure the amount of radiolabeled amino acids in all proteins produced in an in vitro translation reaction including truncated protein products. It is important to separate the radiolabeled protein on a protein gel, and by autoradiography confirm that the product is the proper size and that secondary protein products have not been produced. The most accurate measure of protein

production is to correlate the measure of activity with the measurements of incorporation.

As described above, coupled transcription and translation reactions require the introduction of a DNA template. Further enhancement of in vitro protein translation has been achieved with the use of a vector containing a poly A sequence at one end of the multiple cloning region. A preferred vector also contains an SP6, T7 or T3 RNA polymerase promoter at the opposite end of the multiple cloning region, so that cloning into the vector produces a gene that is flanked by an RNA polymerase promoter at the 5' end and a poly A sequence at the 3' end. Many cloning vectors commercially available contain one or more of the promoters SP6, T7, or T3 as they are widely used for standard in vitro transcription reactions. The vector pSP64 (polyA) is commercially available from Promega Corp., Madison, WI. This vector was used to clone the luciferase gene which was subsequently translated in a coupled transcription and translation reaction using rabbit reticulocyte lysate. The same luciferase gene was cloned into a different vector lacking the poly A sequence and subsequently translated in a coupled transcription and translation reaction. When activity assays were performed on product from these reactions, a significant increase in activity was evident in the reaction which contained the poly A construct.

To study the cotranslational and initial post-translational modifications of proteins, coupled

transcription and translation reactions can be performed in the presence of canine pancreatic microsomal membranes. Signal peptide cleavage, membrane insertion, translocation and core glycosylation are some of the modifications that can be studied. A coupled transcription and translation reaction using a clone of the B-lactamase precursor in the presence of microsomal membranes produced the expected form of the protein showing signal processing had taken place. Likewise, a coupled transcription and translation reaction using a clone of the precursor for  $\alpha$  factor from S. cerevisiae in the presence of microsomal membranes produced the expected processed form of the protein showing glycosylation.

Coupled transcription and translation can be used in any method that requires the in vitro translation of proteins. These methods include in vitro mutagenesis of genes to study structure and function of the resulting proteins. Other methods are the in vitro translation of modified proteins for the purpose of isolating or purifying the protein away from the rest of the reaction and the in vitro translation of protein for the purpose of producing antibodies to that protein. The method of coupled transcription and translation in rabbit reticulocyte lysate or wheat germ extract offers advantages over current methods of in vitro translation in that these systems require less time and yield enhanced levels of protein.

There are also a number of advantages in using such a static coupled transcription and translation reaction over continuous or flow-through reactions. First of all, there is a major difference in the applications for the two systems. The continuous system is for large scale industrial production of proteins whereas static system reactions are suited to the everyday researcher currently doing in vitro translations. Continuous translation is much more expensive to perform, requiring an investment in equipment (a single Amicon unit costs approximately \$2,000) as well as significant amounts of reagents. In particular, the levels of RNA polymerases used to make continuous eukaryotic reactions work is prohibitive for simple research applications (20,000-30,000 units/reaction). Continuous reactions are designed to be performed in relatively large volumes, while static reactions require no extra equipment and only small amounts of reagents, since the reaction volume is typically only on the order of 50  $\mu$ l or less.

The time required for a continuous reaction is significant, anywhere from 24 to 100 hours, whereas static reactions require only 1 to 2 hours to completion. For researchers currently performing in vitro translation, the static system offers significant time savings for most analytical or other research applications by bypassing the in vitro transcription step. Because of the time required both for the set up and for the running of continuous reactions there is no net time savings for a typical researcher, even for

coupled translation and transcription in such a system. Furthermore, with the static coupled transcription and translation system significant increases in protein production are observed when compared to standard in  
5 vitro translation using RNA templates. Further increase in protein production is observed in the static system when the DNA template contains a 3' poly A sequence.

In general, then, the static system makes coupled transcription and translation available to the everyday  
10 researcher, not only because of the cost effectiveness and ease of use, but also because of the significant time savings and increased protein production over other known eukaryotic systems.

#### Example 1

15 Coupled transcription and translation was performed on a DNA construct containing the luciferase gene. The reaction was assayed for the production of luciferase enzyme using the luciferase assay system (Promega, Corp., Madison, WI). Luciferase activity is measured in  
20 Turner light units using a Turner luminometer. Coupled transcription and translation was achieved under the following reaction conditions:

-30-

- 25.0  $\mu$ l standard rabbit reticulocyte lysate  
8.0  $\mu$ l rNTP's (ATP, GTP, UTP, CTP) 2.5 mM each  
1.0  $\mu$ l SP6 RNA polymerase (80 units/ $\mu$ l)  
2.5  $\mu$ l 1xTA buffer\*  
5 1.0  $\mu$ l RNasin (40 units/ $\mu$ l)  
1.0  $\mu$ l pSP64polyA/luc DNA (1  $\mu$ g) circular plasmid DNA  
1.0  $\mu$ l 1 mM amino acids (complete)  
9.5  $\mu$ l H<sub>2</sub>O  
50.0  $\mu$ l

- 10 \* 1xTA buffer consists of 33 mM Tris/acetic acid Ph 7.8,  
65 mM potassium acetate, 10 mM magnesium acetate, 4 mM  
spermidine, and 1 mM DTT.

The reaction was incubated at 30°C for 1.5 hours. After  
1.5 hours a 2.5  $\mu$ l sample from the above reaction was  
15 assayed in the luminometer with a 100x light filter in  
place. The sample produced greater than 30,000 Turner  
light units. Control experiments where either DNA was  
omitted from the reaction or where no 1xTA was included  
produced no Turner light units.

20

Example 2

Coupled transcription and translation was likewise  
performed in a reaction containing wheat germ extract.  
Again, the DNA construct used contained the luciferase  
gene. Coupled transcription and translation was  
25 achieved under the following conditions:

- 25.0  $\mu$ l standard wheat germ extract  
8.0  $\mu$ l rNTP's (ATP, GTP, CTP, UTP) 2.5 mM each  
5.0  $\mu$ l 1xTA buffer  
1.0  $\mu$ l RNasin (40 units/ $\mu$ l)  
30 1.0  $\mu$ l SP6 RNA polymerase (80 units/ $\mu$ l)  
1.0  $\mu$ l pSP64polyA/luc DNA (1  $\mu$ g) circular plasmid DNA  
1.0  $\mu$ l 1 mM amino acids (complete)  
8.0  $\mu$ l H<sub>2</sub>O  
50.0  $\mu$ l

- 35 The reaction was incubated for 1 hour at 30°C. After 1  
hour a 2.5  $\mu$ l sample from the reaction was measured in



-31-

the luminometer with a 100x light filter in place. The sample produced greater than 5,000 Turner light units.

### Example 3

Coupled transcription and translation was also performed with lysates modified during manufacture by the addition of certain components or reagents. During the manufacture of a rabbit reticulocyte lysate an aliquot was set aside and SP6 RNA polymerase was added to the lysate before it was frozen. SP6 was added to the level of 160 units of SP6 per 25  $\mu$ l lysate. The lysate was quick frozen and stored at -70°C until used. Coupled transcription and translation reactions were set up following conditions in example 1 except that no additional SP6 was added. The reactions performed with rabbit reticulocyte lysate that had SP6 RNA polymerase added during manufacture produced protein. Further experiments have shown that the other components (buffers and amino acids, for instance) necessary for coupled transcription and translation can be added to the lysate during its manufacture. The lysate with the components added can be used to produce protein in coupled reactions upon the introduction of DNA templates.

### Example 4

DNA constructs containing genes for a variety of proteins were tested in coupled transcription and translation reactions. The same reaction conditions as

-32-

in example 1 were used except that a radiolabeled amino acid ( $^{35}\text{S}$  methionine) was included in the reaction mix. Methionine was eliminated from the amino acid mix. Reactions were performed using plasmid DNA constructs

5 cloned behind the SP6 or T7 RNA polymerase promoters. pGEMLuc plasmid DNA coding for luciferase was used in coupled transcription and translation reactions and yielded 15.3% incorporation of the radiolabeled amino acid. This compared to a standard in vitro translation

10 using a transcribed RNA template of the luciferase clone which yielded 0.8% incorporation. Samples from these reactions were run on SDS/PAGE gels to confirm that the protein product was the proper size for luciferase. Further experiments were performed using a clones of the

15 B-lactamase gene and the  $\alpha$  factor of S. cerevisiae behind the SP6 promoter. Standard in vitro translation using an RNA template of B-lactamase yielded 3% incorporation of the labeled amino acid whereas a coupled transcription and translation reaction using a

20 DNA template yielded 26.8% incorporation. A standard in vitro translation reaction using an RNA template of the  $\alpha$  factor gene yielded 0.8% incorporation while a coupled transcription and translation reaction using a DNA template yielded 12.4% incorporation. Again, samples of

25 the reactions were run on SDS/PAGE gels to confirm the size of the proteins produced. Other proteins produced from DNA templates in coupled transcription and translation reactions include: TFIID transcription factor, behind the T7 promoter, yielding 5%

-33-

incorporation; Cjun transcription factor, behind T7, yielding 13.4% incorporation; B-galactosidase, behind the SP6 promoter, yielding 28% incorporation; and polyA/luc, a luciferase gene in a poly A vector, behind SP6, yielding 25.9% incorporation.

#### Example 5

Coupled transcription and translation was performed in rabbit reticulocyte lysate using a DNA construct of B-lactamase precursor, in the presence and absence of canine microsomal membranes, to test post-translational modification of proteins. The reactions contained a labeled amino acid in order to visualize the products by autoradiography. Products from the reactions were run on SDS/PAGE gels. The protein product from the reaction without microsomal membranes migrated at about 31.5 kilodaltons (Kd) while the protein product from the reaction containing microsomal membranes migrated at about 28.9 Kd, indicating that the signal sequence had been processed.

In a similar coupled transcription and translation experiment a DNA construct of the  $\alpha$  factor of S. cerevisiae was translated both in the presence and absence of canine microsomal membranes. Again, the radiolabeled products were run on a gel and the results indicated the 18.6 Kd precursor had been processed to a protein migrating at 32.0 Kd, indicating that the  $\alpha$  factor was glycosylated.

-34-  
Example 6

Coupled transcription and translation was performed on DNA produced from a thermocycled amplification process. The only requirement for such a DNA is that the amplified fragment contain an RNA polymerase promoter. The experiment was performed using a DNA fragment amplified from the pGEMLUC plasmid. The resulting fragment contained the sequence for the SP6 RNA polymerase promoter and the sequence for the luciferase gene. When this DNA fragment was introduced into a coupled reaction under conditions similar to example 1, luciferase protein was produced. Other amplified DNA fragments have been translated in coupled reactions including a pGEMEX/gene 10 fragment, behind the T7 promoter, and a B-galactosidase fragment, behind the SP6 promoter.

Although one embodiment of the present invention has been described, it will be apparent to those skilled in the art that various changes and modifications may be made therein without departing from the spirit of the invention or from the scope of the appended claims.

-35-  
CLAIMS

What is claimed is:

1. A method for coupling transcription and translation from DNA in eukaryotic cell-free extract  
5 comprising adding a sufficient amount of a magnesium compound to said extract to raise the magnesium concentration to a level where RNA is transcribed from DNA and RNA translates into protein.
2. The method of Claim 1 wherein said extract is  
10 rabbit reticulocyte lysate.
3. The method of Claim 2 wherein said final magnesium concentration is about 2.5 mM to about 3.5 mM.
4. The method of Claim 2 wherein said final magnesium concentration is about 2.6 mM to about 3.0 mM.
- 15 5. The method of Claim 2 wherein a polyamine is added to said lysate.
6. The method of Claim 5 wherein said polyamine is spermidine.
7. The method of Claim 6 wherein said spermidine  
20 is added to said lysate to a concentration of about 0.2 mM to about 0.4 mM.

8. The method of Claim 2 wherein the potassium concentration of said lysate is adjusted to about 40 mM to about 100 mM.

5           9. The method of Claim 2 including adding a sufficient amount of a ribonuclease inhibitor to said lysate to effectively inactivate endogenous ribonucleases.

10           10. The method of Claim 9 wherein said ribonuclease inhibitor is RNasin.

11. The method of Claim 2 including adding additional components required for coupled transcription and translation to the lysate.

15           12. The method of Claim 11 wherein one of said additional components comprises an RNA polymerase.

13. The method of Claim 12 wherein said polymerase is selected from the group consisting of SP6, T7 and T3 RNA polymerases.

20           14. The method of Claim 2 wherein a DNA template is added to said lysate.

15. The method of Claim 14 wherein said DNA template has a multiple cloning region.

-37-

16. The method of Claim 15 wherein a polymerase promoter sequence is located at one end of said multiple cloning region.

17. The method of Claim 16 wherein said template  
5 has a poly A sequence at the opposite end of said multiple cloning region, so that cloning into said multiple cloning region produces a gene that is flanked by an RNA polymerase promoter at the 5' end and a poly A sequence at the 3' end.

10 18. The method of Claim 16 wherein the polymerase corresponding to said promoter sequence is added to said lysate.

15 19. The method of Claim 14 wherein said DNA template is in a form which is selected from the group consisting of a supercoiled molecule, a covalently closed circular molecule, a linear molecule or a DNA segment made by the process known as a polymerase chain reaction.

20 20. The method of Claim 2 including adding a sufficient amount of ribonucleotide triphosphates to said lysate to allow translation of RNA from DNA.

21. The method of Claim 20 wherein 0.4 mM of each of said ribonucleotide triphosphates are added to said lysate.

22. The method of Claim 1 wherein said extract is wheat germ extract.

23. The method of Claim 22 wherein said final magnesium concentration is about 3.0 mM to about 5.25 mM.

24. The method of Claim 22 wherein said final magnesium concentration is about 4.0 mM to about 4.75 mM.

25. The method of Claim 22 wherein a polyamine is added to said extract.

26. The method of Claim 25 wherein said polyamine is spermidine.

27. The method of Claim 26 wherein said spermidine is added to said extract to a concentration of about 0.2 mM to about 0.9 mM.

28. The method of Claim 22 wherein the potassium concentration of said extract is adjusted to about 50 mM to about 150 mM.

29. The method of Claim 22 including adding a sufficient amount of a ribonuclease inhibitor to said lysate to effectively inactivate endogenous ribonucleases.



-39-

30. The method of Claim 29 wherein said  
ribonuclease inhibitor is RNasin.

31. The method of Claim 22 including adding  
additional components required for coupled transcription  
5 and translation to the lysate.

32. The method of Claim 31 wherein one of said  
additional components comprises an RNA polymerase.

33. The method of Claim 32 wherein said  
polymerase is selected from the group consisting of SP6,  
10 T7 and T3 RNA polymerases.

34. The method of Claim 22 wherein a DNA template  
is added to said extract.

35. The method of Claim 34 wherein said DNA  
template has a multiple cloning region.

15 36. The method of Claim 35 wherein a polymerase  
promoter sequence is located at one end of said multiple  
cloning region.

37. The method of Claim 36 wherein said template  
has a poly A sequence at the opposite end of said  
20 multiple cloning region, so that cloning into said  
multiple cloning region produces a gene that is flanked

-40-

by an RNA polymerase promoter at the 5' end and a poly A sequence at the 3' end.

38. The method of Claim 36 wherein the polymerase corresponding to said promoter sequence is added to said  
5 extract.

39. The method of Claim 34 wherein said DNA template is in a form which is selected from the group consisting of a supercoiled molecule, a covalently closed circular molecule, a linear molecule or a DNA  
10 segment made by the process known as a polymerase chain reaction.

40. The method of Claim 22 including adding a sufficient amount of ribonucleotide triphosphates to said extract to allow translation of RNA from DNA.

15 41. The method of Claim 40 wherein said ribonucleotide triphosphates are added to said extract at the levels of 0.4 mM CTP, 0.4 mM UTP, 0.5 mM GTP and 1.6 mM ATP.

42. A modified eukaryotic cell-free extract  
20 prepared in accordance with Claim 1.

43. A kit comprising a container containing the extract of Claim 42.

- 44 -

44. A modified eukaryotic cell-free extract prepared in accordance with Claim 5 or Claim 25.

45. A kit comprising a container containing the extract of Claim 44.

5           46. A modified eukaryotic cell-free extract prepared in accordance with Claim 12 or Claim 32.

47. A kit comprising a container containing the extract of Claim 46.

10           48. A process for producing protein from a DNA template having a specific polymerase promoter sequence through coupled transcription and translation in a solution, said process comprising the steps of preparing a solution of a standard preparation of a eukaryotic cell-free extract, modifying said extract solution with  
15           sufficient concentrations of ribonucleotide triphosphates, amino acids and the polymerase corresponding to said promoter sequence of said template DNA to support transcription and translation, and adding a sufficient amount of a magnesium compound to said  
20           extract solution to raise the final magnesium concentration to a level where RNA is transcribed from said DNA template and said RNA translates into protein.

-42-

49. The method of Claim 48 wherein the final concentration of magnesium in said extract solution mixture is raised by about 0.5 mM.

50. The method of Claim 48 wherein said extract is rabbit reticulocyte lysate.

51. The method of Claim 50 wherein said final magnesium concentration is about 2.5 mM to about 3.5 mM.

52. The method of Claim 50 wherein said final magnesium concentration is about 2.6 mM to about 3.0 mM.

53. The method of Claim 50 wherein a polyamine is added to said solution.

54. The method of Claim 53 wherein said polyamine is spermidine.

55. The method of Claim 54 wherein said spermidine is added to said solution to a concentration of about 0.2 mM to about 0.4 mM.

56. The method of Claim 50 wherein the potassium concentration of said solution is adjusted to about 40 mM to about 100 mM.

57. The method of Claim 50 including adding a sufficient amount of a ribonuclease inhibitor to said

-43-

solution to effectively inactivate endogenous ribonucleases.

58. The method of Claim 57 wherein said ribonuclease inhibitor is RNasin.

5 59. The method of Claim 50 including adding additional components required for coupled transcription and translation to the solution.

60. The method of Claim 59 wherein one of said additional components comprises an RNA polymerase.

10 61. The method of Claim 60 wherein said polymerase is selected from the group consisting of SP6, T7 and T3 RNA polymerases.

62. The method of Claim 50 wherein said DNA template is in a form which is selected from the group  
15 consisting of a supercoiled molecule, a covalently closed circular molecule, a linear molecule or a DNA segment made by the process known as a polymerase chain reaction.

63. The method of Claim 50 wherein said DNA  
20 template has a multiple cloning region.

- 44 -

64. The method of Claim 63 wherein a polymerase promoter sequence is located at one end of said multiple cloning region.

5 65. The method of Claim 64 wherein said template has a poly A sequence at the opposite end of said multiple cloning region, so that cloning into said multiple cloning region produces a gene that is flanked by an RNA polymerase promoter at the 5' end and a poly A sequence at the 3' end.

10 66. The method of Claim 50 wherein 0.4 mM of each of said ribonucleotide triphosphates are added to said solution.

67. The method of Claim 48 wherein said extract is wheat germ extract.

15 68. The method of Claim 67 wherein said final magnesium concentration is about 3.0 mM to about 5.25 mM.

69. The method of Claim 68 wherein said final magnesium concentration is about 4.0 mM to about 4.75  
20 mM.

70. The method of Claim 67 wherein a polyamine is added to said solution.

71. The method of Claim 70 wherein said polyamine is spermidine.

72. The method of Claim 71 wherein said spermidine is added to said solution to a concentration  
5 of about 0.2 mM to about 0.9 mM.

73. The method of Claim 67 wherein the potassium concentration of said solution is adjusted to about 50 mM to about 150 mM.

74. The method of Claim 67 including adding a  
10 sufficient amount of a ribonuclease inhibitor to said solution to effectively inactivate endogenous ribonucleases.

75. The method of Claim 74 wherein said ribonuclease inhibitor is RNasin.

15 76. The method of Claim 67 including adding additional components required for coupled transcription and translation to the solution.

77. The method of Claim 76 wherein one of said additional components comprises an RNA polymerase.

20 78. The method of Claim 77 wherein said polymerase is selected from the group consisting of SP6, T7 and T3 RNA polymerases.

79. The method of Claim 67 wherein said DNA  
template is in a form which is selected from the group  
consisting of a supercoiled molecule, a covalently  
closed circular molecule, a linear molecule or a DNA  
5 segment made by the process known as a polymerase chain  
reaction.

80. The method of Claim 67 wherein said DNA  
template has a multiple cloning region.

81. The method of Claim 80 wherein a polymerase  
10 promoter sequence is located at one end of said multiple  
cloning region.

82. The method of Claim 81 wherein said template  
has a poly A sequence at the opposite end of said  
multiple cloning region, so that cloning into said  
15 multiple cloning region produces a gene that is flanked  
by an RNA polymerase promoter at the 5' end and a poly A  
sequence at the 3' end.

83. The method of Claim 67 wherein said  
ribonucleotide triphosphates are added to said solution  
20 at the levels of 0.4 mM CTP, 0.4 mM UTP, 0.5 mM GTP and  
1.6 mM ATP.



## AMENDED CLAIMS

[received by the International Bureau on 18 March 1993 (18.03.93);  
original claims 11,31,59 and 76 cancelled; original claims 1,12,  
20,32,48,60,66 and 77 amended; new claims 84 and 85 added;  
other claims unchanged (9 pages)]

1. A method for coupling transcription and translation from DNA in eukaryotic cell-free extract  
5 comprising adding a sufficient amount of a magnesium salt to said extract to raise the magnesium concentration to a level where RNA is transcribed from DNA and RNA translates into protein.
2. The method of Claim 1 wherein said extract is  
10 rabbit reticulocyte lysate.
3. The method of Claim 2 wherein said final magnesium concentration is about 2.5 mM to about 3.5 mM.
4. The method of Claim 2 wherein said final magnesium concentration is about 2.6 mM to about 3.0 mM.
- 15 5. The method of Claim 2 wherein a polyamine is added to said lysate.
6. The method of Claim 5 wherein said polyamine is spermidine.
7. The method of Claim 6 wherein said spermidine  
20 is added to said lysate to a concentration of about 0.2 mM to about 0.4 mM.

8. The method of Claim 2 wherein the potassium concentration of said lysate is adjusted to about 40 mM to about 100 mM.

5           9. The method of Claim 2 including adding a sufficient amount of a ribonuclease inhibitor to said lysate to effectively inactivate endogenous ribonucleases.

10           10. The method of Claim 9 wherein said ribonuclease inhibitor is RNasin.

12. The method of Claim 2 wherein an RNA polymerase is added to the lysate.

15           13. The method of Claim 12 wherein said polymerase is selected from the group consisting of SP6, T7 and T3 RNA polymerases.

14. The method of Claim 2 wherein a DNA template is added to said lysate.

15. The method of Claim 14 wherein said DNA template has a multiple cloning region.

16. The method of Claim 15 wherein a polymerase promoter sequence is located at one end of said multiple cloning region.

5 17. The method of Claim 16 wherein said template has a poly A sequence at the opposite end of said multiple cloning region, so that cloning into said multiple cloning region produces a gene that is flanked by an RNA polymerase promoter at the 5' end and a poly A sequence at the 3' end.

10 18. The method of Claim 16 wherein the polymerase corresponding to said promoter sequence is added to said lysate.

15 19. The method of Claim 14 wherein said DNA template is in a form which is selected from the group consisting of a supercoiled molecule, a covalently closed circular molecule, a linear molecule or a DNA segment made by the process known as a polymerase chain reaction.

20 20. The method of Claim 2 including adding a sufficient amount of natural ribonucleotide triphosphates to said lysate to allow translation of RNA from DNA.

25 21. The method of Claim 20 wherein 0.4 mM of each of said ribonucleotide triphosphates are added to said lysate.

30. The method of Claim 29 wherein said ribonuclease inhibitor is RNasin.

32. The method of Claim 22 wherein an RNA polymerase is added to the extract.

5           33. The method of Claim 32 wherein said polymerase is selected from the group consisting of SP6, T7 and T3 RNA polymerases.

34. The method of Claim 22 wherein a DNA template is added to said extract.

10           35. The method of Claim 34 wherein said DNA template has a multiple cloning region.

36. The method of Claim 35 wherein a polymerase promoter sequence is located at one end of said multiple cloning region.

15           37. The method of Claim 36 wherein said template has a poly A sequence at the opposite end of said multiple cloning region, so that cloning into said multiple cloning region produces a gene that is flanked

44. A modified eukaryotic cell-free extract prepared in accordance with Claim 5 or Claim 25.

45. A kit comprising a container containing the extract of Claim 44.

5           46. A modified eukaryotic cell-free extract prepared in accordance with Claim 12 or Claim 32.

47. A kit comprising a container containing the extract of Claim 46.

10           48. A process for producing protein from a DNA template having a specific polymerase promoter sequence through coupled transcription and translation in a solution, said process comprising the steps of preparing a solution of a standard preparation of a eukaryotic cell-free extract containing said DNA template,  
15           modifying said extract solution with sufficient concentrations of natural ribonucleotide triphosphates, amino acids and the polymerase corresponding to said promoter sequence of said template DNA to support transcription and translation, and adding a sufficient  
20           amount of a magnesium salt to said extract solution to raise the final magnesium concentration to a level where RNA is transcribed from said DNA template and said RNA translates into protein.

solution to effectively inactivate endogenous ribonucleases.

58. The method of Claim 57 wherein said ribonuclease inhibitor is RNasin.

5 60. The method of Claim 50 wherein an RNA polymerase is added to the lysate.

61. The method of Claim 60 wherein said polymerase is selected from the group consisting of SP6, T7 and T3 RNA polymerases.

10 62. The method of Claim 50 wherein said DNA template is in a form which is selected from the group consisting of a supercoiled molecule, a covalently closed circular molecule, a linear molecule or a DNA segment made by the process known as a polymerase chain  
15 reaction.

63. The method of Claim 50 wherein said DNA template has a multiple cloning region.

64. The method of Claim 63 wherein a polymerase promoter sequence is located at one end of said multiple cloning region.

5 65. The method of Claim 64 wherein said template has a poly A sequence at the opposite end of said multiple cloning region, so that cloning into said multiple cloning region produces a gene that is flanked by an RNA polymerase promoter at the 5' end and a poly A sequence at the 3' end.

10 66. The method of Claim 50 wherein 0.4 mM of each of said natural ribonucleotide triphosphates are added to said solution.

67. The method of Claim 48 wherein said extract is wheat germ extract.

15 68. The method of Claim 67 wherein said final magnesium concentration is about 3.0 mM to about 5.25 mM.

20 69. The method of Claim 68 wherein said final magnesium concentration is about 4.0 mM to about 4.75 mM.

70. The method of Claim 67 wherein a polyamine is added to said solution.

71. The method of Claim 70 wherein said polyamine is spermidine.

72. The method of Claim 71 wherein said spermidine is added to said solution to a concentration  
5 of about 0.2 mM to about 0.9 mM.

73. The method of Claim 67 wherein the potassium concentration of said solution is adjusted to about 50 mM to about 150 mM.

74. The method of Claim 67 including adding a  
10 sufficient amount of a ribonuclease inhibitor to said solution to effectively inactivate endogenous ribonucleases.

75. The method of Claim 74 wherein said ribonuclease inhibitor is RNasin.

15 77. The method of Claim 67 wherein an RNA polymerase is added to the extract.

78. The method of Claim 77 wherein said polymerase is selected from the group consisting of SP6, T7 and T3 RNA polymerases.

20 79. The method of Claim 67 wherein said DNA template is in a form which is selected from the group consisting of a supercoiled molecule, a covalently closed circular molecule, a linear molecule or a DNA segment made by the process known as a polymerase chain  
25 reaction.



80. The method of Claim 67 wherein said DNA template has a multiple cloning region.

81. The method of Claim 80 wherein a polymerase promoter sequence is located at one end of said multiple cloning region.

82. The method of Claim 81 wherein said template has a poly A sequence at the opposite end of said multiple cloning region, so that cloning into said multiple cloning region produces a gene that is flanked by an RNA polymerase promoter at the 5' end and a poly A sequence at the 3' end.

83. The method of Claim 67 wherein said ribonucleotide triphosphates are added to said solution at the levels of 0.4 mM CTP, 0.4 mM UTP, 0.5 mM GTP and 1.6 mM ATP.

84. The method of Claim 1 wherein said transcription and translation are coupled in a batch reaction.

85. The method of Claim 48 wherein said transcription and translation are coupled in a batch reaction.

## STATEMENT UNDER ARTICLE 19

In response to the Notification of Transmittal of the International Search Report mailed January 12, 1993, the following amendment is presented in accordance with the Regulations under the Patent Cooperation Treaty, Rule 46. Pursuant to 46.1, and presuming this communication is received before the technical preparations for international publication are completed, applicant requests that the enclosed replacement sheets be substituted in the above application. The amendments contained in the sheets, which are described more fully below, are intended only to bring the application into better conformance with the application as currently prosecuted before the United States Patent & Trademark Office.

Claims 1, 12, 20, 32, 48, 60, 66 and 77 are amended. Claims 84 and 85 are new, while claims 11, 31, 59 and 76 are canceled. These changes necessitate the submission of replacement sheets for pages 35, 36, 37, 39, 41, 43, 44, 45, and 46 of the original application. The amendments to the claims bring them into closer conformance with the application being prosecuted in the United States Patent & Trademark Office.

In Claims 1 and 48, "compound" has been replaced with ---salt---.

In Claims 12, 32, 60 and 77 the dependency has been changed and an RNA polymerase is added to the lysate.

In Claims 20, 48 and 66 the term ---natural--- has been inserted before "ribonucleotide".

In Claim 48 the eukaryotic cell-free extract is now described as specifically containing a DNA template.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US92/08518

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C12P 21/00

US CL : 435/68.1

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/68.1, 69.1, 370.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

DIALOG, BIOSIS, MEDLINE, APS

SEARCH TERMS: COUPLED, TRANSLATION, TRANSCRIPTION, IN VITRO

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<u>X</u> Y	WO, A, 91/02076 (BARANOV ET AL.) 21 FEBRUARY 1991, SEE ENTIRE DOCUMENT.	1, 2, 5- 10, 22, 25 -30, 32-40, 42-48, 50, 53-58, 60 -65, 67, 70 <del>-75, 77-82</del> 3-4, 21, 23-24, 41, 49, 51-52, 66, 68-69, 83-85
Y	J. BIOCHEM. VOLUME 82, ISSUED 1977, SUZUKI, "EFFECT OF KCL, MAGNESIUM ACETATE AND SPERMINE ON THE RATION OF A TO B GLOBIN CHAINS SYNTHESIZED", PAGES 251-260, SEE ENTIRE DOCUMENT.	3-4, 23-24, 41, 49, 51-52, 66, 68-69 AND 83
Y	VIROLOGY VOLUME 73, ISSUED 1976, BREINDL ET AL., "STUDIES ON THE IN VITRO TRANSCRIPTION AND TRANSLATION OF VESICULAR STOMATITIS VIRUS mRNA" PAGES 106-118, SEE ENTIRE DOCUMENT.	1-83

☒ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A		document defining the general state of the art which is not considered to be part of particular relevance
* E		earlier document published on or after the international filing date
* L		document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
* O		document referring to an oral disclosure, use, exhibition or other means
* P		document published prior to the international filing date but later than the priority date claimed
	* X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
	* Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
	* A	document member of the same patent family

Date of the actual completion of the international search

06 January 1993

Date of mailing of the international search report

12 JAN 1993

Name and mailing address of the ISA/  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

DAVID B. SCHMICKEL

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-4202

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/US92/08518

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PROC. NATL. ACAD. SCI. USA. VOLUME 70, NO. 8, ISSUED AUGUST 1973, ROBERTS ET AL. "EFFICIENT TRANSLATION OF TOBACCO MOSAIC VIRUS RNA AND RABBIT GLOBIN 9S RNA IN A CELL-FREE SYSTEM FROM COMERCIAL WHEAT GERM", PAGES 2330-2334, SEE ENTIRE DOCUMENT.	1-83
Y	SCIENCE, VOLUME 242, ISSUED 25 NOVEMBER 1988, SPIRIN ET AL., "A CONTINUOUS CELL-FREE TRANSLATION SYSTEM CAPABLE OF PRODUCING POLYPEPTIDES IN HIGH YIELD", PAGES 1162-1164, SEE ENTIRE DOCUMENT.	1-83
Y	NUCLEIC ACIDS RES., VOLUME 17, NUMBER 11, ISSUED JUNE 1989, RYABOVA ET AL., "PREPATATIVE SYNTHESIS IN A CONTINUOUS CELL-FREE SYSTEM FROM RABBIT RETICULOCYTES" PAGE 4412, SEE FIGURE 1.	1-83
Y	GENE, VOLUME 84, ISSUED 1989, BARANOV ET AL., "GENE EXPRESSION IN A CELL-FREE SYSTEM ON THE PREPARATIVE SCALE", PAGES 463-466, SEE FIGURE 1, AND ENTIRE DOCUMENT.	1-83